

Intrinsic transverse momentum in hard processes¹

P.J. Mulders

*Department of Physics and Astronomy, Free University
 and
 National Institute for Nuclear Physics and High Energy Physics (NIKHEF)
 P.O. Box 41882, NL-1009 DB Amsterdam, The Netherlands*

Abstract


In this contribution I discuss transverse momentum dependent quark distributions and their operator structure. Some of the experimental signatures in particular in Drell-Yan scattering are given.

1 Introduction

In hard processes one becomes sensitive to intrinsic transverse momentum of the quarks if at least three external momenta are present. In Drell-Yan (DY) scattering ($AB \rightarrow \mu^+ \mu^- X$) one has the momenta of the incoming hadrons A and B and the hard time-like momentum of the lepton-pair q (with $q^2 = Q^2$). Other examples are 1-particle inclusive leptonproduction ($\ell H \rightarrow \ell' h X$) and two-hadron production in $e^+ e^-$ annihilation.


In a field-theoretic approach one can calculate the hadronic tensor $W_{\mu\nu}$ for DY scattering by making use of a diagrammatic expansion in which soft parts allow inclusion of hadrons in the field-theoretical framework of QCD. If $Q^2 \rightarrow \infty$ only a limited number of diagrams survives, each of these diagrams with characteristic behavior. Some explicit examples are shown below. In particular the tree level diagram in the first box and the diagrams with a gluon attached to the soft part (one shown in the second box) will be discussed in more detail.

¹ Talk presented at the Symposium "Deep inelastic scattering off polarized targets: theory meets experiment", Sep. 1-5, 1997, Zeuthen



2 Soft parts

The soft parts that appear in the diagrammatic expansion are the quark-quark correlation function,




$$p^2 \sim p \cdot P \sim P^2 = M^2 \ll Q^2$$

given by

$$\Phi_{ij}(p; P, S) = \frac{1}{(2\pi)^4} \int d^4x e^{ip \cdot x} \langle P, S | \bar{\psi}_j(0) \psi_i(x) | P, S \rangle, \quad (1)$$

and the quark-quark-gluon correlation function



$$k^2 \sim k \cdot P \sim k_1^2 \sim k_1 \cdot P \sim k \cdot k_1 \sim P^2 = M^2 \ll Q^2$$

given by

$$\Phi_{Aij}^\alpha(k, k_1; P, S) = \frac{1}{(2\pi)^4} \int d^4x d^4y e^{ik_1 \cdot (x-y) + ik \cdot y} \langle P, S | \bar{\psi}_j(0) g A_T^\alpha(y) \psi_i(x) | P, S \rangle. \quad (2)$$

The large scale Q in the scattering process leads in a natural way to the use of lightlike vectors for which $n_+^2 = n_-^2 = 0$ and $n_+ \cdot n_- = 1$. For the DY process:

$$\left. \begin{aligned} q^2 &= Q^2 \\ P_A^2 &= M_A^2 \\ P_B^2 &= M_B^2 \\ 2 P_A \cdot q &= Q^2/x_A \\ 2 P_B \cdot q &= Q^2/x_B \end{aligned} \right\} \longleftrightarrow \begin{cases} P_B = \frac{Q}{x_B \sqrt{2}} n_- + \frac{x_B M_B^2}{Q \sqrt{2}} n_+ \\ q = \frac{Q}{\sqrt{2}} n_- + \frac{Q}{\sqrt{2}} n_+ + q_T \\ P_A = \frac{x_A M_A^2}{Q \sqrt{2}} n_- + \frac{Q}{x_A \sqrt{2}} n_+ \end{cases}$$

Using the lightlike vectors to define lightcone components $p^\pm = p \cdot n_\mp$ one sees that the relevant quantities entering in the hard process for the parton graph in the first box above are $\int dp^- \Phi$ and $\int dk^+ \bar{\Phi}$. The structure of these quantities, their various Dirac projections, and a possible way of estimating their magnitude has been discussed in the contribution of Jakob [1]. The relevant part in $\int dp^- \Phi$,

$$\begin{aligned} \Phi_{ij}(x, \mathbf{p}_T) &= \int dp^- \Phi_{ij}(p; P, S) \\ &= \int \frac{d\xi^- d^2 \xi_T}{(2\pi)^3} e^{ip \cdot \xi} \langle P, S | \bar{\psi}_j(0) \psi_i(\xi) | P, S \rangle \Big|_{\xi^+=0}, \end{aligned} \quad (3)$$

which is a light-front correlation function, in leading order in $1/Q$ given by [2, 3]

$$\begin{aligned} \Phi(x, \mathbf{p}_T) &= \frac{1}{2} \left\{ f_1 \not{n}_+ + \lambda g_{1L} \gamma_5 \not{n}_+ + g_{1T} \frac{(\mathbf{p}_T \cdot \mathbf{S}_T)}{M} \gamma_5 \not{n}_+ + h_{1T} \frac{[\not{\mathbf{S}}_T, \not{n}_+] \gamma_5}{2} \right. \\ &\quad \left. + \lambda h_{1L}^\perp \frac{[\not{\mathbf{p}}_T, \not{n}_+] \gamma_5}{2} + h_{1T}^\perp \frac{(\mathbf{p}_T \cdot \mathbf{S}_T)}{M} \frac{[\not{\mathbf{p}}_T, \not{n}_+] \gamma_5}{2} \right\} + \mathcal{O}\left(\frac{M}{P^+}\right), \end{aligned} \quad (4)$$

where $p^+ = x P^+$ and the functions $f_1 = f_1(x, \mathbf{p}_T^2)$ etc. The polarization vector is also expanded with the help of the lightlike vectors,

$$S = -\lambda \frac{M}{2P^+} n_- + \lambda \frac{P^+}{M} n_+ + S_T. \quad (5)$$

and satisfies $\lambda^2 + \mathbf{S}_T^2 = 1$ (for a pure state). The functions appearing in $\Phi(x, \mathbf{p}_T)$ can be projected out with specific Dirac matrices. All functions in the leading part can be interpreted as densities involving chirally left/right or transverse spin projections of good quark fields, $\psi_+ \equiv (\gamma^- \gamma^+ / 2) \psi$.

In inclusive lepton-hadron scattering or $\mathbf{q}_T$ -integrated DY scattering one does not need the $\mathbf{p}_T$ -dependent quark distributions discussed above, but just the $\mathbf{p}_T$ -averaged results, i.e.

$$\begin{aligned} \Phi_{ij}(x) &= \int dp^- d^2 \mathbf{p}_T \Phi_{ij}(p; P, S) \\ &= \int \frac{d\xi^-}{2\pi} e^{ip \cdot \xi} \langle P, S | \bar{\psi}_j(0) \psi_i(\xi) | P, S \rangle \Big|_{LC}, \end{aligned} \quad (6)$$

where 'LC' denotes $\xi^+ = \xi_T = 0$, which implies $\xi^2 = 0$, i.e. one deals with a lightcone correlation function. In leading order one has

$$\Phi(x) = \frac{1}{2} \left\{ f_1(x) \not{n}_+ + \lambda g_1(x) \gamma_5 \not{n}_+ + h_1(x) \frac{[\not{\mathcal{S}}_T, \not{n}_+] \gamma_5}{2} \right\} + \mathcal{O}\left(\frac{M}{P^+}\right), \quad (7)$$

where $f_1(x) = \int d^2 \mathbf{p}_T f_1(x, \mathbf{p}_T)$, etc. For asymmetries in 1-particle inclusive lepton-hadron scattering or in DY scattering one also needs the $\mathbf{p}_T$ -weighted results,

$$\begin{aligned} \Phi_{\partial ij}^\alpha(x) &= \int dp^- d^2 \mathbf{p}_T p_T^\alpha \Phi_{ij}(p; P, S) \\ &= \int \frac{d\xi^-}{2\pi} e^{ip \cdot \xi} \langle P, S | \bar{\psi}_j(0) i \partial^\alpha \psi_i(\xi) | P, S \rangle \Big|_{LC}, \end{aligned} \quad (8)$$

for which one in leading order has

$$\frac{1}{M} \Phi_{\partial}^\alpha(x) = \frac{1}{2} \left\{ g_{1T}^{(1)}(x) S_T^\alpha \gamma_5 \not{n}_+ + \lambda h_{1L}^{(1)}(x) \frac{[\not{n}_+, \gamma^\alpha] \gamma_5}{2} \right\} + \mathcal{O}\left(\frac{M}{P^+}\right), \quad (9)$$

with e.g. $g_{1T}^{(1)}(x) \equiv \int d^2 \mathbf{p}_T (\mathbf{p}_T^2 / 2M^2) g_{1T}(x, \mathbf{p}_T)$.

Including subleading order for the $\mathbf{p}_T$ -averaged functions we just quote the result

$$\Phi(x) = \frac{1}{2} \left\{ f_1(x) \not{n}_+ + \lambda g_1(x) \gamma_5 \not{n}_+ + h_1(x) \frac{[\not{\mathcal{S}}_T, \not{n}_+] \gamma_5}{2} \right\} \quad (10)$$

$$+ \frac{M}{2P^+} \left\{ e(x) + g_T(x) \gamma_5 \not{\mathcal{S}}_T + \lambda h_L(x) \frac{[\not{n}_+, \not{n}_-] \gamma_5}{2} \right\} + \mathcal{O}\left(\frac{M}{P^+}\right)^2. \quad (11)$$

The factor M/P^+ leads in a calculation to a suppression factor M/Q , hence the functions e , g_T and h_L appear at subleading order. These functions do not have a simple density interpretation. From the most general amplitude expansion for Φ_{ij} , however, constrained by hermiticity, parity and time reversal symmetry one obtains relations between the twist-three functions and the $\mathbf{p}_T$ -dependent distribution functions [4, 5],

$$\underbrace{g_T(x) - g_1(x)}_{g_2(x)} = \frac{d}{dx} g_{1T}^{(1)}, \quad (12)$$

$$\underbrace{h_L(x) - h_1(x)}_{\frac{1}{2} h_2(x)} = -\frac{d}{dx} h_{1L}^{\perp(1)}. \quad (13)$$

3 Gluons in the diagrammatic calculation

The soft parts discussed in the previous section and the analogous results for antiquarks are sufficient to calculate the parton diagram (first box in the introduction). Turning to the second box one finds diagrams with a gluon attached to the soft part involving the matrix elements of Eq. 2. These contributions are important at subleading order [6]. The nonlocality up to $\mathcal{O}(1/Q)$ can be restricted to a lightlike nonlocality involving $\bar{\psi}(0) A_T^\alpha(\xi) \psi(\xi)$ with $\xi^+ = \xi_T = 0$. At least, this is true in the absence of so-called gluonic poles [7], which is the case when gluonic background matrix elements having a nonlocality of the type $\bar{\psi}(0) A_T^\alpha(\eta^-) \psi(\xi^-)$ vanish.

To ensure gauge invariance one looks for the combination $p_T^\alpha + g A_T^\alpha$, i.e. for quark-quark correlation functions containing the covariant derivative $iD_T^\alpha = i\partial_T^\alpha + g A_T^\alpha$

$$\begin{aligned} \Phi_{Aij}^\alpha(x) &= \int \frac{d\xi^-}{2\pi} e^{ip \cdot \xi} \langle P, S | \bar{\psi}_j(0) g A_T^\alpha(\xi) \psi_i(\xi) | P, S \rangle \Big|_{LC} \\ &= \Phi_{Dij}^\alpha(x) - \Phi_{\partial ij}^\alpha(x). \end{aligned} \quad (14)$$

In the next step one employs the equations of motion, $(i\gamma^\mu D_\mu - m)\psi = 0$, to relate Φ_D to Φ . In this way one obtains

$$\begin{aligned} \Phi_D^\alpha(x) &= \frac{M}{2} \left\{ \left(x g_T - \frac{m}{M} h_1 \right) S_T^\alpha \gamma_5 \not{p}_+ + \lambda \left(x h_L - \frac{m}{M} g_1 \right) \frac{[\gamma^\alpha, \not{p}_+] \gamma_5}{4} \right. \\ &\quad \left. + \left(x e - \frac{m}{M} f_1 \right) \frac{[\not{p}_+, \gamma^\alpha]}{2} \right\}. \end{aligned} \quad (15)$$

In the diagrammatic calculation up to $\mathcal{O}(1/Q)$ the parton graph and the gluon graphs ensure electromagnetic gauge invariance [6]. The difference between Φ_D^α and Φ_∂^α gives the truly interaction dependent part,

$$\Phi_A^\alpha(x) = \frac{M}{2} \left\{ x \tilde{g}_T(x) S_T^\alpha \gamma_5 \not{p}_+ + \lambda x \tilde{h}_L(x) \frac{[\gamma^\alpha, \not{p}_+] \gamma_5}{4} + x \tilde{e}(x) \frac{[\not{p}_+, \gamma^\alpha]}{2} \right\}, \quad (16)$$

with

$$x \tilde{g}_T = x g_T - g_{1T}^{(1)} - \frac{m}{M} h_1, \quad (17)$$

$$x \tilde{h}_L = x h_L - h_{1L}^{\perp(1)} - \frac{m}{M} g_1, \quad (18)$$

$$x \tilde{e} = x e - \frac{m}{M} f_1. \quad (19)$$

As an aside we note that one obtains the Wandzura-Wilczek relation [8] for g_T and a similar relation [6] for h_L ,

$$g_T(x) = \int_x^1 dy \frac{g_1(y)}{y} + \frac{m}{M} \left[\frac{h_1(x)}{x} - \int_x^1 dy \frac{h_1(y)}{y^2} \right], \quad (20)$$

$$h_L(x) = 2x \int_x^1 dy \frac{h_1(y)}{y^2} + \frac{m}{M} \left[\frac{g_1(x)}{x} - 2x \int_x^1 dy \frac{g_1(y)}{y^3} \right], \quad (21)$$

by using the relations in Eqs 12 and 13 and assuming $\Phi_A^\alpha = 0$.

Included in the diagrammatic expansion are also quark-quark-gluon correlations involving A^+ gluons. These longitudinal A^+ gluons cause problems in the M/P^+ expansion. Their contributions, however, can be collected into the definition of the distribution functions. One obtains for instance


$$f_1(x) = \int \frac{d\xi^-}{4\pi} e^{ip \cdot \xi} \langle P, S | \bar{\psi}(0) \gamma^+ \mathcal{P} e^{-ig \int d\eta^\mu A_\mu(\eta)} \psi(\xi) | P, S \rangle \Big|_{LC}, \quad (22)$$

rendering the definition of f_1 color gauge invariant. In cases such as inclusive deep inelastic lepton-hadron scattering all complications can simply be avoided by working in the lightcone gauge ($A^+ = 0$).

Including longitudinal gluons and transverse momenta we get

$$\Phi_{ij}(x, \mathbf{p}_T) = \int \frac{d\xi^- d^2 \boldsymbol{\xi}_T}{2(2\pi)^3} e^{ip \cdot \xi} \langle P, S | \bar{\psi}_j(0) \mathcal{U}(0, \infty) \mathcal{U}(\infty, \xi^-) \psi_i(\xi) | P, S \rangle \Big|_{\xi^+ = 0}, \quad (23)$$

where link operators at fixed transverse positions,



appear of the form

$$\mathcal{U}(\xi, \eta) = \mathcal{P} e^{-ig \int_\xi^\eta d\zeta^- A^+(\zeta)} \Big|_{\xi_T = \eta_T = \zeta_T}. \quad (24)$$

The integrated results become


$$\Phi_{ij}(x) = \int \frac{d\xi^-}{4\pi} e^{ip \cdot \xi} \langle P, S | \bar{\psi}_j(0) \mathcal{U}(0, \xi) \psi_i(\xi) | P, S \rangle \Big|_{LC}, \quad (25)$$

$$\begin{aligned} \Phi_{\partial ij}^\alpha(x) = \int \frac{d\xi^-}{4\pi} e^{ip \cdot \xi} \Big\{ & \langle P, S | \bar{\psi}_j(0) \mathcal{U}(0, \xi) i D_T^\alpha \psi_i(\xi) | P, S \rangle \Big|_{LC} \\ & - \langle P, S | \bar{\psi}_j(0) \mathcal{U}(0, \infty) \int_{\infty}^{\xi^-} d\eta^- \mathcal{U}(\infty, \eta) g F^{+\alpha}(\eta) \mathcal{U}(\eta, \xi) \psi_i(\xi) | P, S \rangle \Big|_{LC} \Big\}. \end{aligned} \quad (26)$$

We note that in $A^+ = 0$ gauge: $F^{+\alpha} = \partial^+ A_T^\alpha$ from which one obtains $A_T^\alpha(\xi) = \int_{\infty}^{\xi^-} d\eta^- F^{+\alpha}(\eta)$.

4 Cross sections

We start with the for our purposes relevant quantities for the DY process given in the next box. This includes the invariants x_A , x_B and y as well as a conveniently chosen cartesian set of vectors, which also serves to define azimuthal angles.



lepton plane (cm)

Cartesian set

$$\begin{aligned} T^\mu &= q^\mu \\ Z^\mu &= x_A P_A^\mu - x_B P_B^\mu \\ X^\mu &= q^\mu - x_A P_A^\mu - x_B P_B^\mu \end{aligned}$$

Note: $T^2 = Q^2$, $Z^2 = -Q^2$, $X^2 = -Q_T^2$

$$AB \longrightarrow \ell \bar{\ell} X$$

$$x_A = \frac{Q^2}{2P_A \cdot q}$$

$$x_B = \frac{Q^2}{2P_B \cdot q}$$

$$y = \frac{1}{2} (1 + \cos \theta)$$

$$\approx \frac{k \cdot P_A}{q \cdot P_A} \approx \frac{k' \cdot P_B}{q \cdot P_B}$$

We will be interested in particular in weighted cross sections involving azimuthal angles and define

$$\langle W \rangle_{\underbrace{P_A P_B}_{\uparrow}} \equiv \int d\phi^\ell d^2 \mathbf{q}_T \underbrace{W}_{\uparrow} \frac{d\sigma_{P_A P_B}}{dx_A dx_B dy d\phi^\ell d^2 \mathbf{q}_T}, \quad (27)$$

$$(O, L, T) \quad W(Q_T, \phi_h^\ell, \phi_{S_A}^\ell, \phi_{S_B}^\ell) \quad (28)$$

where O , L , T indicate unpolarized, longitudinally polarized ($\lambda = 1$) or transversely polarized ($|\mathbf{S}_T| = 1$) hadrons. The azimuthal angles $\phi_h^\ell = \phi_h - \phi^\ell$ are defined w.r.t. the lepton plane.

One has e.g. for the simplest unpolarized $\mathbf{q}_T$ -averaged cross section the well-known result

$$\langle 1 \rangle_{OO} = \frac{\pi\alpha^2}{3Q^2} (1 + \cos^2 \theta) \sum_{a,\bar{a}} e_a^2 f_1^a(x_A) f_1^{\bar{a}}(x_B), \quad (29)$$

where we have included the summation over quarks and antiquarks and the flavor indices on the distribution functions.

We start with a discussion of *spin asymmetries in polarized $\mathbf{q}_T$ -integrated DY scattering*, where one finds the following nonvanishing asymmetries,

$$\langle 1 \rangle_{LL} = \frac{\pi\alpha^2}{3Q^2} \lambda_A \lambda_B (1 + \cos^2 \theta) \sum_{a,\bar{a}} e_a^2 g_1^a(x_A) g_1^{\bar{a}}(x_B), \quad (30)$$

$$\langle \cos(\phi_{S_A}^\ell + \phi_{S_B}^\ell) \rangle_{TT} = \frac{\pi\alpha^2}{6Q^2} \lambda_A \lambda_B (1 - \cos^2 \theta) \sum_{a,\bar{a}} e_a^2 h_1^a(x_A) h_1^{\bar{a}}(x_B), \quad (31)$$

$$\begin{aligned} \langle \cos(\phi_{S_B}^\ell) \rangle_{LT} = \frac{\pi\alpha^2}{3Q^2} |\mathbf{S}_{AT}| |\mathbf{S}_{BT}| \sin 2\theta \sum_{a,\bar{a}} e_a^2 \left\{ \frac{M_A x_A}{Q} (h_L^a + \tilde{h}_L^a)(x_A) h_1^{\bar{a}}(x_B) \right. \\ \left. + \frac{M_B x_B}{Q} g_1^a(x_A) (g_T^{\bar{a}} + \tilde{g}_T^{\bar{a}})(x_B) \right\} \end{aligned} \quad (32)$$

These asymmetries have been given in Ref. [6] disregarding $\mathbf{k}_T$ -dependence of the distribution functions. In that case one has for the twist-three functions $\tilde{g}_T = g_T$ and $\tilde{h}_L = h_L$, which simplifies the above LT asymmetry. However, taking into account the transverse momentum dependence, both the interaction-dependent (tilde) functions and the full twist-three functions including a twist two part appear in the LT asymmetry [9].

As a next example, we consider two specific *azimuthal asymmetries in polarized DY scattering (at leading order)*, namely

$$\left\langle \frac{Q_T}{M_B} \cos(\phi_h^\ell - \phi_{S_B}^\ell) \right\rangle_{LT} = \frac{\pi\alpha^2}{3Q^2} \lambda_A |\mathbf{S}_{BT}| (1 + \cos^2 \theta) \sum_{a,\bar{a}} e_a^2 g_1^a(x_A) g_{1T}^{(1)\bar{a}}(x_B), \quad (33)$$

$$\left\langle \frac{Q_T}{M_A} \cos(\phi_h^\ell + \phi_{S_B}^\ell) \right\rangle_{LT} = \frac{\pi\alpha^2}{3Q^2} \lambda_A |\mathbf{S}_{BT}| (1 - \cos^2 \theta) \sum_{a,\bar{a}} e_a^2 h_{1L}^{\perp(1)a}(x_A) h_1^{\bar{a}}(x_B) \quad (34)$$

discussed in Ref. [3]. Together with the previous subleading LT asymmetries these allow tests of the relations in Eqs 12 and 13. We note that the constraints on the parametrization of the quark-quark correlation functions, notably time reversal invariance only allow double spin asymmetries in DY scattering.

5 Time reversal symmetry

Non-applicability of time reversal symmetry (T) allows a richer structure for Φ . This is the case in, for instance, the corresponding definitions for fragmentation functions [10, 5], where one cannot use T as a symmetry because the final state hadron is not a plane

wave state. But also for distribution functions it has been argued that time reversal odd quantities may appear, e.g. through soft factorization breaking mechanisms, the chiral structure of nucleons or gluonic poles [11, 12, 7]. In the quark-quark correlation function then the following structures are allowed

$$\Phi(x, \mathbf{p}_T) \Big|_{T\text{-odd}} = \frac{1}{2} \left\{ f_{1T}^\perp(x, \mathbf{k}_T) \frac{\epsilon_{\mu\nu\rho\sigma} \gamma^\mu n_+^\nu k_T^\rho S_T^\sigma}{M} + h_1^\perp(x, \mathbf{k}_T) \frac{i[\not{p}_T, \not{k}_+]}{2M} \right\}. \quad (35)$$

Integrating over $\mathbf{p}_T$ this T-odd part vanishes, but the $\mathbf{p}_T$ -weighted result is nonvanishing,

$$\Phi_\partial^\alpha(x) \Big|_{T\text{-odd}} = \frac{M}{2} \left\{ -f_{1T}^{\perp(1)}(x) \epsilon^\alpha_{\mu\nu\rho} \gamma^\mu n_-^\nu S_T^\rho - h_1^{\perp(1)}(x) \frac{i[\gamma^\alpha, \not{k}_+]}{2} \right\}. \quad (36)$$

We note that, as in the T-even situation, the twist-three T-odd distribution functions showing up in the order (M/P^+) part of Φ ,

$$\Phi(x) \Big|_{T\text{-odd}} = \frac{M}{2P^+} \left\{ f_T(x) \epsilon_T^{\rho\sigma} S_{T\rho} \gamma_\sigma - \lambda e_L(x) i\gamma_5 + h(x) \frac{i[\not{k}_+, \not{k}_-]}{2} \right\}, \quad (37)$$

can be related to the $\mathbf{k}_T^2$ -moments in Φ_∂^α ,

$$h(x) = -\frac{d}{dx} h_1^{\perp(1)}, \quad (38)$$

$$f_T(x) = -\frac{d}{dx} f_{1T}^{\perp(1)}. \quad (39)$$

At $\mathcal{O}(1/Q)$ one again needs the quark-quark-gluon correlation functions and the use of the equations of motion to rewrite

$$\Phi_D^\alpha(x) \Big|_{T\text{-odd}} = \frac{M}{2} \left\{ x f_T(x) \epsilon^\alpha_{\mu\nu\rho} \gamma^\mu n_-^\nu S_T^\rho + x h(x) \frac{i[\gamma^\alpha, \not{k}_+]}{4} \right\}. \quad (40)$$

The interaction-dependent parts can again be isolated,

$$\Phi_A^\alpha(x) \Big|_{T\text{-odd}} = \frac{M}{2} \left\{ x \tilde{f}_T(x) \epsilon^\alpha_{\mu\nu\rho} \gamma^\mu n_-^\nu S_T^\rho + x \tilde{h}(x) \frac{i[\gamma^\alpha, \not{k}_+]}{4} \right\}, \quad (41)$$

with

$$x \tilde{f}_T = x f_T + f_{1T}^{\perp(1)}, \quad (42)$$

$$x \tilde{h} = x h + 2 h_1^{\perp(1)}. \quad (43)$$

The equivalence of the Wandzura-Wilczek parts, obtained by combining $\Phi_A^\alpha = 0$ with Eqs 38 and 39 is simply $f_T(x) = h(x) = 0$.

As an example of T-odd distributions we mention the following *azimuthal asymmetry for unpolarized leptonproduction (leading twist)*

$$\left\langle \frac{Q_T^2}{MM_h} \cos(2\phi_h^\ell) \right\rangle_{OOO} = \frac{16\pi\alpha^2 s}{Q^4} (1-y) \sum_{a,\bar{a}} e_a^2 x_B h_1^{\perp(1)a}(x_B) H_1^{\perp(1)a}(z_h), \quad (44)$$

which involves besides the T-odd distribution function $h_1^{\perp(1)}$, depending on $x_B = Q^2/2P \cdot q$, the equivalent fragmentation function $H_1^{\perp(1)}$ depending on $z_h = P \cdot P_h/P \cdot q$. Examples of single spin asymmetries in the DY process are given in the talk of Boer [7].

6 Conclusions

I have reviewed the systematic parametrization of quark-quark and quark-quark-gluon correlation functions which appear as the soft parts in the diagrammatic expansion of hard processes such as DY scattering. In particular, I have emphasized the role of transverse momenta of quarks, accounting for which is essential in processes like DY scattering or 1-particle inclusive lepton-hadron scattering. The effects of transverse momenta of quarks show up, for instance, in azimuthal asymmetries. While at leading order the relevant functions in the soft parts of hard processes are just the quark densities, one sees at subleading order the QCD dynamics at work inside the hadrons.

This work is part of the scientific program of the foundation for Fundamental Research on Matter (FOM) and the Dutch Organization for Scientific Research (NWO).

References

- [1] R. Jakob, P.J. Mulders and J. Rodrigues, contribution in these proceedings; see also [hep-ph/9704335](#).
- [2] J.P. Ralston and D.E. Soper, *Nucl. Phys.* **B172** (1980) 445.
- [3] R.D. Tangerman and P.J. Mulders, *Phys. Rev.* **D51** (1995) 3357.
- [4] A.P. Bukhvostov, E.A. Kuraev, L.N. Lipatov, *SP JETP* **60** (1984) 22.
- [5] P.J. Mulders and R.D. Tangerman, *Nucl. Phys.* **B461** (1996) 197 and Erratum, *Nucl. Phys.* **B484** (1997) 538.
- [6] R.L. Jaffe and X. Ji, *Nucl. Phys.* **B375** (1992) 527.
- [7] D. Boer, P.J. Mulders and O.V. Teryaev, contribution in these proceedings and refs therein; see also [hep-ph/9710223](#).
- [8] S. Wandzura and F. Wilczek, *Phys. Lett.* **B72** (1977) 195.

- [9] R.D. Tangerman and P.J. Mulders, [hep-ph/9408305](#).
- [10] R.L. Jaffe and X. Ji, *Phys. Rev. Lett.* **71** (1993) 2547.
- [11] D. Sivers, *Phys. Rev.* **D41** (1990) 83 and *Phys. Rev.* **D43** (1991) 261;
J. Collins, *Nucl. Phys.* **B396** (1993) 161.
- [12] M. Anselmino, M. Boglione and F. Murgia, *Phys. Lett.* **B362** (1995) 164;
M. Anselmino, A. Drago and F. Murgia, [hep-ph/9703303](#);
M. Anselmino, contribution in these proceedings, [hep-ph/9710330](#).